

Chronic Obstructive Pulmonary Diseases: Journal of the COPD Foundation



COPD9USA Session Summary

COPD Overlap Syndromes: Asthma and Beyond

Allison A. Lambert, MD, MHS¹ and Mark T. Dransfield, MD²

This article serves as a CME-available enduring material summary of the following COPD9USA presentations:

- “COPD and Asthma” Presenter: Prescott Woodruff, MD, MPH
- “COPD and Lung Cancer” Presenter: William Bulman, MD
- “COPD and Bronchiectasis” Presenter: Jeremy Clain, MD
- “COPD and Interstitial Lung Disease” Presenter: George Washko, MD

Abbreviations: chronic obstructive pulmonary disease, **COPD**; asthma-COPD overlap syndrome, **ACOS**; computerized tomography, **CT**; Global Initiative for Asthma, **GINA**; Global initiative for chronic Obstructive Lung Disease, **GOLD**; The Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points, **ECLIPSE**; intensive care unit, **ICU**; lung volume reduction surgery, **LVRS**; high resolution computerized tomography, **HRCT**; combined pulmonary fibrosis and emphysema, **CPFE**

Citation: Lambert AA, Dransfield MT. COPD overlap syndromes: Asthma and beyond. *Chronic Obstr Pulm Dis (Miami)*. 2016;3(1):459-465. doi: <http://dx.doi.org/10.15326/jcopdf.3.1.2015.0176>

1 Department of Internal Medicine Division of Pulmonary and Critical Care Medicine, Johns Hopkins University, Baltimore, Maryland

2 Division of Pulmonary, Allergy and Critical Care and Lung Health Center, University of Alabama at Birmingham and the Birmingham Veterans Administration Medical Center

Address correspondence to:

Allison Lambert, MD, MHS
1830 E. Monument Street
5th Floor
Division of Pulmonary and Critical Care Medicine
Baltimore, MD 21205
Email: alamber5@jhmi.edu

Keywords:

COPD; overlap; asthma; lung cancer; bronchiectasis; interstitial lung disease

Introduction

Chronic obstructive pulmonary disease (COPD) affects 15 million Americans and became the third leading cause of death in the United States in 2011.¹ Comorbidities are increasingly recognized to play an important role in the management and outcomes of patients with COPD.^{2,3} COPD phenotypes have been proposed to facilitate patient classification for

prognostic, therapeutic, and research purposes.⁴ Proposed phenotypes have been expanded beyond subtypes of COPD to include overlapping pulmonary disease.⁵ Overlapping diagnoses with COPD complicate the diagnosis and management of this disease but may also pose an opportunity to identify distinct and modifiable pathophysiology. We summarize here 4 of these overlapping syndromes that were reviewed at the 2015 COPD9USA conference.

COPD and Asthma

The overlap of COPD with asthma is referred to as asthma-COPD overlap syndrome (ACOS). ACOS represents a clinical diagnosis and should be considered among patients with persistent airflow limitation and a combination of features typically associated with both asthma and COPD.⁶ Examples of features more typical of asthma than COPD that may be present include: childhood symptoms, significant day to day variability of symptoms, known triggers for symptoms, personal or family history of atopy, or more pronounced reversibility of airflow obstruction. Despite these characteristics, classic COPD features may still be present, such as hyperinflation or computerized tomography (CT)-defined emphysema. Identification of patients with

ACOS is critical because outcomes for patients with ACOS may be worse than those for patients with COPD or asthma alone;⁷ as a result, subspecialty referral is often required to ensure appropriate diagnosis and treatment.

Although the Global Initiative for Asthma (GINA) and the Global initiative for chronic Obstructive Lung Disease (GOLD) recommend a clinical definition of ACOS,⁶ characterizing ACOS by these clinical features may not represent the distinguishing underlying pathophysiology and may inappropriately categorize patients. Therefore, distinguishing ACOS from singular COPD may be more appropriately performed by utilizing biological features that may reflect the underlying disease pathology and the response to therapy.⁸

Eosinophilia may represent the appropriate surrogate to distinguish these clinical diseases. A subset of patients with COPD are known to exhibit eosinophilic, rather than neutrophilic, inflammation.⁹ The Evaluation of COPD Longitudinally to Identify Predictive Surrogate End-points (ECLIPSE) cohort observed 37% of participants to have persistently elevated blood eosinophil counts ($\geq 2\%$) over 3 years and reported that these patients demonstrated distinct characteristics, including older age, male gender, fewer symptoms and less airflow obstruction.¹⁰ Importantly, ECLIPSE found that blood and sputum eosinophilia were correlated. Sputum eosinophilia has been found to correlate with responsiveness to both systemic^{11,12} and inhaled^{13,14} corticosteroids among COPD patients. Additional targeted therapy beyond corticosteroids has been suggested for eosinophilic COPD patients with the medication benralizumab, a monoclonal antibody that inhibits interleukin-5.¹⁵

In summary, ACOS represents a distinct disease process with both unique pathophysiology and responsiveness to therapy. As opposed to traditional clinically-driven phenotypes, ACOS may be better defined by biological tools, such as blood or sputum eosinophil count. Use of these biologic markers may more accurately characterize the underlying pathophysiology and predict targeted therapeutic responsiveness.

COPD and Lung Cancer


Lung cancer is the leading cancer cause of death among both men and women in the United States¹⁶ and kills more people annually than colorectal, breast, prostate and pancreatic cancers combined.¹⁷ More

men are diagnosed with lung cancer each year but the incidence among women has been rising, related to changing smoking patterns over the last century. Approximately 42 million American adults currently smoke cigarettes,¹⁸ 12-15 million have COPD,^{1,19} and approximately 400,000 have lung cancer.²⁰ Figure 1 illustrates the prevalence of these overlapping diagnoses.

Approximately 50%-80% of patients with lung cancer had pre-existing COPD prior to their cancer diagnosis.²¹⁻²³ Comorbid COPD and lung cancer is prevalent due to the shared risk factor of tobacco use however, the risk for lung cancer among patients with COPD exceeds expected rates based upon age or extent of tobacco exposure.²⁴⁻²⁶ Specifically, COPD is independently associated with a 2.5-fold increase in the odds of lung cancer, after adjusting for age, gender, and extent of tobacco use.²⁷ COPD patients with CT-defined emphysema have a nearly 4-fold independent increase in the odds of lung cancer.²⁸ This independently elevated risk for lung cancer among COPD patients is therefore not likely driven by tobacco use or age alone, but rather other processes such as chronic inflammation, amplified carcinogenic effects from tobacco exposure, impaired mucociliary clearance or shared genetic susceptibility.²⁹

Not only do patients with COPD experience increased risk for lung cancer compared to their smoking, age-matched peers, but they also suffer worse prognosis from

Figure 1. Prevalence of Smoking, COPD and Lung Cancer



Graphic illustrates prevalence of cigarette smoking (42.1 million in 2013), COPD diagnosis (12 million in 2010) and lung diagnosis cancer (408,808 people in 2012) among adults in the United States.

lung cancer with approximately doubled mortality.^{30,31} GOLD guidelines suggest that patients with comorbid COPD and lung cancer should receive the same treatment for each condition as would be recommended for patients with either diagnosis alone³²; however these recommendations are often limited by the severity of underlying respiratory disease including surgical, chemotherapeutic and radioablative thresholds.^{29,33} An increasing number of studies are now focused on expanding application of these therapies beyond traditional thresholds or identifying less invasive or aggressive treatment regimens that may still afford benefit in those with advanced chronic pulmonary disease.^{34,35} Curative surgical resection for early stage lung cancer has been an area of focus because it offers the greatest mortality benefit for patients with early disease,³⁶ may preserve lung function in those with limited reserve and may provide benefits similar to lung volume reduction surgery (LVRS) in select candidates. LVRS showed the greatest benefit for patients with upper lobe predominant emphysema and poor exercise capacity following pulmonary rehabilitation³⁷; therefore these COPD patients who have resectable lung cancer in an upper lobe may derive dual benefit from surgery.

Given the known poor prognosis for lung cancer among COPD patients and the many limitations to providing standard therapies for these patients, prevention and screening are critical components of care. Aggressive tobacco cessation efforts³⁸ and targeted screening chest CT scans³⁹ are imperative in the care of COPD patients. Importantly, COPD patients in the CT arm of the National Lung Screening Trial did not experience overdiagnosis with CT screening but rather a shift towards a more favorable stage at the time of cancer diagnosis.⁴⁰

COPD and Bronchiectasis

Bronchiectasis, a suppurative lung disease with radiographically defined airway dilation, is common among patients with advanced COPD and is distinct from airway wall thickening which is a common finding in obstructive lung diseases.⁴¹ Comorbid bronchiectasis has been reported in 28%-58% of COPD patients with moderate to severe airflow obstruction⁴²⁻⁴⁴ with increasing prevalence as degree of airflow obstruction increases.⁴⁵ These reports may underestimate the true burden of bronchiectasis because bronchiectasis is a radiographically-defined diagnosis that exhibits clinical overlap with COPD; therefore, COPD patients may not

be appropriately evaluated for comorbid bronchiectasis. As the performance of chest CTs increases for the purposes of lung cancer screening, bronchiectasis will likely become increasingly recognized.

Despite this clinical overlap, recognition of bronchiectasis among patients with COPD is important because its presence is associated with worse outcomes. Specifically, outpatients with comorbid bronchiectasis and COPD suffer more frequent respiratory exacerbations, higher rates of potentially pathogenic microorganisms isolated in their sputum, and increased mortality.⁴⁶ Intubated intensive care unit (ICU) patients with overlapping bronchiectasis and COPD experience prolonged courses of mechanical ventilation, extended ICU and hospital length-of-stay, and increased rates of ventilator-associated pneumonia.⁴⁷

In addition to prognostication and risk stratification, diagnosis of bronchiectasis may also carry therapeutic relevance regarding the approach to microorganisms isolated in sputum, the role for bronchial hygiene, and the possibility of targeted pulmonary resection. In non-cystic fibrosis-related bronchiectasis, colonization with *Pseudomonas aeruginosa* has been associated both with disease progression, defined by forced expiratory volume in 1 second decline,⁴⁸ and mortality.⁴⁹ Anti-pseudomonal treatment is recommended for non-cystic fibrosis patients with evidence of chronic colonization with *Pseudomonas aeruginosa*⁵⁰ and may also have a role for those with frequent exacerbations.⁵¹ Application of these findings to COPD patients with overlapping bronchiectasis and frequent exacerbations has been proposed but is not yet approved⁵²; therefore, individual practitioners may consider use of these therapies in a case-by-case fashion, without Federal Drug Administration approval.

In summary, bronchiectasis is prevalent among patients with advanced COPD and may represent a distinct phenotype that warrants targeted therapy.

COPD and Interstitial Lung Disease

Emphysema and fibrosis are radiographically and physiologically defined features that individually represent the extremes of lung disease but can coexist. Emphysema is characterized by a loss of lung tissue, decreased lung recoil and airflow obstruction; whereas fibrosis is characterized by a gain of lung tissue, increased lung recoil and airflow restriction. The overlap of these 2 conditions, likely related to the common exposure of tobacco smoke or other noxious

particles or gasses, results in pseudo-normalization of spirometry and lung volumes and an isolated, severely reduced diffusion capacity.⁵³ Resting and exertional hypoxemia are common,⁵⁴ with 1 study reporting 80% of combined pulmonary fibrosis and emphysema (CPFE) patients required supplemental oxygen.⁵⁵ The overlap of COPD and interstitial lung disease is often overlooked when imaging, specifically high resolution CT (HRCT), is not available to demonstrate the combination of radiographic pulmonary emphysema and fibrosis. CPFE was first recognized in 1990 but the prevalence of these overlapping diseases has become increasingly apparent as the performance of HRCTs increases.^{54,56} Despite this, targeted, specific therapy still does not exist.

Presently, research is focused upon identifying radiographic precursors of CPFE to identify those at risk for disease earlier, at which time interventions may afford greater benefit.^{57,58} COPD Genetic Epidemiology is a prospective cohort study that has obtained regular CT imaging on current and former smokers over time. At the time of enrollment into this study, 8% of participants demonstrated interstitial lung abnormalities.⁵⁹ These participants demonstrated reduced CT lung volume, less emphysema, greater symptoms, and reduced 6 minute walk distance as compared to their smoking counterparts without interstitial lung abnormalities on CT.⁶⁰ Shared genetic polymorphisms are being investigated to allow the identification of patients at risk for the overlap of emphysema with fibrosis, thereby enabling focused screening and therapy. The Muc5B promoter polymorphism, highly associated with idiopathic pulmonary fibrosis, is now recognized to be associated with interstitial lung disease in the general population.⁶¹

In summary, CPFE is a prevalent condition associated with worse outcomes in the COPD population. Radiographic and genetic methods are being developed to identify patients at risk for this manifestation of COPD. These tools will additionally allow the identification of optimal subpopulations for clinical therapeutic trials.

Conclusions

In conclusion, COPD is a prevalent and morbid condition, complicated by comorbid diseases and overlapping pulmonary syndromes. Better characterization of these overlapping syndromes may afford enhanced diagnostic methods, prognostication, and ultimately therapeutic intervention and outcomes.

Declaration of Interest

Dr. Lambert has served as a consultant for GLG Consulting and has received grants from the National Institutes of Health. Dr. Dransfield has received grants from GlaxoSmithKline and Forest and has been a contracted researcher for Boehringer Ingelheim, GlaxoSmithKline, Pulmonx, PheumRx, Otsuka, Pearl, Forest and AstraZeneca. Dr. Woodruff has served as an advisor for Janssen, Neostem and Genentech. He has been a consultant for Roche, Novartis and AstraZeneca and has received a grant from Genentech. Dr. Washko has served as a consultant for GlaxoSmithKline and his spouse is employed by Merck.

NOTE: To complete the CME post test for this article, refer to the original online version of the article at:
www.journal.copdfoundation.org

References

- Centers for Disease Control and Prevention. Chronic Obstructive Pulmonary Disease. Centers for Disease Control and Prevention website. <http://www.cdc.gov/copd/index.html>. Published March 2015. Accessed December 20, 2015.
- Barnes PJ, Celli BR. Systemic manifestations and comorbidities of COPD. *Eur Respir J*. 2009;33 (5):1165-1185. doi: <http://dx.doi.org/10.1183/09031936.00128008>
- Fabbri LM, Luppi F, Beghe B, Rabe KF. Complex chronic comorbidities of COPD. *Eur Respir J*. 2008;31(1):204-212. doi: <http://dx.doi.org/10.1183/09031936.00114307>
- Han MK, Agusti A, Calverley PM, et al. Chronic obstructive pulmonary disease phenotypes: the future of COPD. *Am J Respir Crit Care Med*. 2010;182(5):598-604. doi: <http://dx.doi.org/10.1164/rccm.200912-1843CC>
- Rennard SI, Locantore N, Delafont B, et al. Identification of five chronic obstructive pulmonary disease subgroups with different prognoses in the ECLIPSE cohort using cluster analysis. *Annals Am Thoracic Soc*. 2015;12(3):303-12. doi: <http://dx.doi.org/10.1513/AnnalsATS.201403-125OC>
- Global Initiative for Asthma. Diagnosis of Diseases of Chronic Airflow Limitation: Asthma COPD and Asthma - COPD (ACOS). Global initiative for chronic Lung Disease website. <http://www.goldcopd.org/uploads/users/files/AsthmaCOPDOverlap.pdf> Published 2014. Accessed December 20, 2015.
- Alshabanat A, Zafari Z, Albanyan O, Dairi M, FitzGerald JM. Asthma and COPD Overlap Syndrome (ACOS): A Systematic Review and Meta Analysis. *PloS One*. 2015;10:e0136065. doi: <http://dx.doi.org/10.1371/journal.pone.0136065>
- Pavord ID, Shaw DE, Gibson PG, Taylor DR. Inflammometry to assess airway diseases. *Lancet*. 2008;372(9643):1017-1019. doi: [http://dx.doi.org/10.1016/S0140-6736\(08\)61421-X](http://dx.doi.org/10.1016/S0140-6736(08)61421-X)
- Saha S, Brightling CE. Eosinophilic airway inflammation in COPD. *Int J Chron Obstruct Pulmon Dis*. 2006;1(1):39-47.
- Singh D, Kolsum U, Brightling CE, et al. Eosinophilic inflammation in COPD: Prevalence and clinical characteristics. *Eur Respir J*. 2014;44(6):1697-700. doi: <http://dx.doi.org/10.1183/09031936.00162414>
- Brightling CE, Monteiro W, Ward R, et al. Sputum eosinophilia and short-term response to prednisolone in chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet*. 2000;356(9240):1480-1485. doi: [http://dx.doi.org/10.1016/S0140-6736\(00\)02872-5](http://dx.doi.org/10.1016/S0140-6736(00)02872-5)
- Siva R, Green RH, Brightling CE, et al. Eosinophilic airway inflammation and exacerbations of COPD: a randomised controlled trial. *Eur Respir J*. 2007;29(5):906-913. doi: <http://dx.doi.org/10.1183/09031936.00146306>
- Brightling CE, McKenna S, Hargadon B, et al. Sputum eosinophilia and the short term response to inhaled mometasone in chronic obstructive pulmonary disease. *Thorax*. 2005;60:193-198. doi: <http://dx.doi.org/10.1136/thx.2004.032516>
- Leigh R, Pizzichini MM, Morris MM, Maltais F, Hargreave FE, Pizzichini E. Stable COPD: predicting benefit from high-dose inhaled corticosteroid treatment. *Eur Respir J*. 2006;27:964-971. doi: <http://dx.doi.org/10.1183/09031936.06.00072105>
- Brightling CE, Bleecker ER, Panettieri RA, Jr, et al. Benralizumab for chronic obstructive pulmonary disease and sputum eosinophilia: a randomised, double-blind, placebo-controlled, phase 2a study. *Lancet Respir Med*. 2014;2(11):891-901. doi: [http://dx.doi.org/10.1016/S2213-2600\(14\)70187-0](http://dx.doi.org/10.1016/S2213-2600(14)70187-0)
- Centers for Disease Control and Prevention. Lung Cancer. Centers for Disease Control and Prevention website. <http://www.cdc.gov/cancer/lung/> Published September 2015. Accessed December 20, 2015.
- American Lung Association. Lung Cancer Fact Sheet: Estimated Cancer Deaths by Site, 2015. American Lung Association website. <http://ala1-old.pub30.convio.net/lung-disease/lung-cancer/resources/facts-figures/lung-cancer-fact-sheet.html> Published 2015. Accessed December 20, 2015
- Centers for Disease Control and Prevention. Smoking and Tobacco Use: Current Cigarette Smoking Among Adults in the United States. Centers for Disease Control and Prevention website. http://www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking/ Updated 2015. Accessed December 20, 2015.
- National Institutes of Health. Research Portfolio Online Reporting Tools (RePORT): COPD. Updated March 29, 2015. Accessed December 20, 2015.
- National Institutes of Health-National Cancer Institute. SEER Stat Fact Sheets: Lung and Bronchus Cancer. National Cancer Institute website. <http://seer.cancer.gov/statfacts/html/lungb.html> Accessed December 20, 2015.
- Young RP, Hopkins RJ, Christmas T, Black PN, Metcalf P, Gamble GD. COPD prevalence is increased in lung cancer, independent of age, sex and smoking history. *Eur Respir J*. 2009;34:380-386. doi: <http://dx.doi.org/10.1183/09031936.00144208>
- Loganathan RS, Stover DE, Shi W, Venkatraman E. Prevalence of COPD in women compared to men around the time of diagnosis of primary lung cancer. *Chest*. 2006;129(5):1305-1312. doi: <http://dx.doi.org/10.1378/chest.129.5.1305>
- Congleton J, Muers MF. The incidence of airflow obstruction in bronchial carcinoma, its relation to breathlessness, and response to bronchodilator therapy. *Respir Med*. 1995;89(4):291-296. doi: [http://dx.doi.org/10.1016/0954-6111\(95\)90090-X](http://dx.doi.org/10.1016/0954-6111(95)90090-X)
- Tockman MS, Anthonisen NR, Wright EC, Donithan MG. Airways obstruction and the risk for lung cancer. *Ann Intern Med*. 1987;106(4):512-518. doi: <http://dx.doi.org/10.7326/0003-4819-106-4-512>
- Skillrud DM, Offord KP, Miller RD. Higher risk of lung cancer in chronic obstructive pulmonary disease. A prospective, matched, controlled study. *Ann Intern Med*. 1986;105(4):503-537. doi: <http://dx.doi.org/10.7326/0003-4819-105-4-503>
- de Torres JP, Marin JM, Casanova C, et al. Lung cancer in patients with chronic obstructive pulmonary disease-- incidence and predicting factors. *Am J Respir Crit Care Med*. 2011;184(8):913-919. doi: <http://dx.doi.org/10.1164/rccm.201103-0430OC>
- Koshiol J, Rotunno M, Consonni D, et al. Chronic obstructive pulmonary disease and altered risk of lung cancer in a population-based case-control study. *PloS One*. 2009;4:e7380. doi: <http://dx.doi.org/10.1371/journal.pone.0007380>

28. Li Y, Swensen SJ, Karabekmez LG, et al. Effect of emphysema on lung cancer risk in smokers: a computed tomography-based assessment. *Cancer Prev Res*. 2011;4:43-50. doi: <http://dx.doi.org/10.1158/1940-6207.CAPR-10-0151>
29. Raviv S, Hawkins KA, DeCamp MM, Jr., Kalhan R. Lung cancer in chronic obstructive pulmonary disease: enhancing surgical options and outcomes. *Am J Respir Crit Care Med*. 2011;183(9):1138-46. doi: <http://dx.doi.org/10.1164/rccm.201008-1274CI>
30. Aldrich MC, Munro HM, Mumma M, et al. Chronic obstructive pulmonary disease and subsequent overall and lung cancer mortality in low-income adults. *PLoS One*. 2015;10:e0121805. doi: <http://dx.doi.org/10.1371/journal.pone.0121805>
31. Kiri VA, Soriano J, Visick G, Fabbri L. Recent trends in lung cancer and its association with COPD: an analysis using the UK GP Research Database. *Prim Care Respir J*. doi: <http://dx.doi.org/10.4104/pcrj.2009.00048>
32. Global initiative for chronic Lung Disease (GOLD). Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease. GOLD website. http://www.goldcopd.org/uploads/users/files/GOLD_Report_2015_Sept2.pdf Updated 2015. Accessed December 20, 2015.
33. Bolliger CT, Perruchoud AP. Functional evaluation of the lung resection candidate. *Eur Respir J*. 1998;11(1):198-212. <http://dx.doi.org/10.1183/09031936.98.11010198>
34. Izquierdo JL, Resano P, El Hachem A, Graziani D, Almonacid C, Sanchez IM. Impact of COPD in patients with lung cancer and advanced disease treated with chemotherapy and/or tyrosine kinase inhibitors. *Int J Chron Obstr Pulmon Dis*. 2014;9(1):1053-1058. doi: <http://dx.doi.org/10.2147/COPD.S68766>
35. Licker MJ, Widikker I, Robert J, et al. Operative mortality and respiratory complications after lung resection for cancer: impact of chronic obstructive pulmonary disease and time trends. *Ann Thorac Surg*. 2006;81:1830-1837. doi: <http://dx.doi.org/10.1016/j.athoracsur.2005.11.048>
36. Van Schil PE. Surgery for non-small cell lung cancer. *Lung Cancer*. 2001;34 (Suppl 2):S127-132. doi: [http://dx.doi.org/10.1016/S0169-5002\(01\)00354-3](http://dx.doi.org/10.1016/S0169-5002(01)00354-3)
37. Fishman A, Martinez F, Naunheim K, et al. A randomized trial comparing lung-volume-reduction surgery with medical therapy for severe emphysema. *New Eng J Med*. 2003;348:2059-2073. doi: <http://dx.doi.org/10.1056/NEJMoa030287>
38. Tashkin DP, Murray RP. Smoking cessation in chronic obstructive pulmonary disease. *Respir Med*. 2009;103:963-974. doi: <http://dx.doi.org/10.1016/j.rmed.2009.02.013>
39. National Lung Screening Trial Research Team, Aberle DR, Adams AM, et al. Reduced lung-cancer mortality with low-dose computed tomographic screening. *New Eng J Med*. 2011;365:395-409. doi: <http://dx.doi.org/10.1056/NEJMoa1102873>
40. Young RP, Duan F, Chiles C, et al. Airflow limitation and histology shift in the National Lung Screening Trial. The NLST-ACRIN Cohort Substudy. *Am J Respir Crit Care Med*. 2015;192(9):1060-1067. doi: <http://dx.doi.org/10.1164/rccm.201505-0894OC>
41. McShane PJ, Naureckas ET, Tino G, Strek ME. Non-cystic fibrosis bronchiectasis. *Am J Respir Crit Care Med*. 2013;188(6):647-56. doi: <http://dx.doi.org/10.1164/rccm.201303-0411CI>
42. Martinez-Garcia MA, Soler-Cataluna JJ, Donat Sanz Y, et al. Factors associated with bronchiectasis in patients with COPD. *Chest*. 2011;140(5):1130-1137. doi: <http://dx.doi.org/10.1378/chest.10-1758>
43. Patel IS, Vlahos I, Wilkinson TM, et al. Bronchiectasis, exacerbation indices, and inflammation in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2004;170(4):400-407. doi: <http://dx.doi.org/10.1164/rccm.200305-648OC>
44. O'Brien C, Guest PJ, Hill SL, Stockley RA. Physiological and radiological characterisation of patients diagnosed with chronic obstructive pulmonary disease in primary care. *Thorax*. 2000;55(8):635-642. doi: <http://dx.doi.org/10.1136/thorax.55.8.635>
45. Agusti A, Calverley PM, Celli B, et al. Characterisation of COPD heterogeneity in the ECLIPSE cohort. *Respir Res*. 2010;11:122.
46. Martinez-Garcia MA, de la Rosa Carrillo D, Soler-Cataluna JJ, et al. Prognostic value of bronchiectasis in patients with moderate-to-severe chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2013;187(8):823-831. doi: <http://dx.doi.org/10.1164/rccm.201208-1518OC>
47. Gursel G. Does coexistence with bronchiectasis influence intensive care unit outcome in patients with chronic obstructive pulmonary disease? *Heart Lung*. 2006;35:58-65.
48. Martinez-Garcia MA, Soler-Cataluna JJ, Perpina-Tordera M, Roman-Sanchez P, Soriano J. Factors associated with lung function decline in adult patients with stable non-cystic fibrosis bronchiectasis. *Chest*. 2007;132(5):1565-1572. doi: <http://dx.doi.org/10.1378/chest.07-0490>
49. Loebinge MR, Wells AU, Hansell DM, et al. Mortality in bronchiectasis: a long-term study assessing the factors influencing survival. *Eur Respir J*. 2009;34:843-849. doi: <http://dx.doi.org/10.1183/09031936.00003709>
50. Pasteur MC, Bilton D, Hill AT for British Thoracic Society Bronchiectasis non CF Guideline Group. British Thoracic Society guideline for non-CF bronchiectasis. *Thorax*. 2010;65 (Suppl 1):i1-58. doi: <http://dx.doi.org/10.1136/thx.2010.136119>
51. Quon BS, Goss CH, Ramsey BW. Inhaled antibiotics for lower airway infections. *Ann Thoracic Soc*. 2014;11(3):425-434. doi: <http://dx.doi.org/10.1513/AnnalsATS.201311-395FR>
52. Novosad SA, Barker AF. Chronic obstructive pulmonary disease and bronchiectasis. *Curr Opin Pulm Med*. 2013;19:133-139. doi: <http://dx.doi.org/10.1097/MCP.0b013e32835d8312>
53. Jankowich MD, Rounds SI. Combined pulmonary fibrosis and emphysema syndrome: a review. *Chest*. 2012;141(1):222-231. doi: <http://dx.doi.org/10.1378/chest.11-1062>
54. Cottin V, Nunes H, Brillet PY, et al. Combined pulmonary fibrosis and emphysema: a distinct underrecognised entity. *Eur Respir J*. 2005;26(4):586-593. doi: <http://dx.doi.org/10.1183/09031936.05.00021005>
55. Jankowich MD, Rounds S. Combined pulmonary fibrosis and emphysema alters physiology but has similar mortality to pulmonary fibrosis without emphysema. *Lung*. 2010;188(5):365-373. doi: <http://dx.doi.org/10.1007/s00408-010-9251-6>

-
56. Wiggins J, Strickland B, Turner-Warwick M. Combined cryptogenic fibrosing alveolitis and emphysema: the value of high resolution computed tomography in assessment. *Respir Med.* 1990;84(5):365-369.
doi: [http://dx.doi.org/10.1016/S0954-6111\(08\)80070-4](http://dx.doi.org/10.1016/S0954-6111(08)80070-4)
-
57. Tsushima K, Sone S, Yoshikawa S, Yokoyama T, Suzuki T, Kubo K. The radiological patterns of interstitial change at an early phase: over a 4-year follow-up. *Respir Med.* 2010;104(11):1712-1721.
doi: <http://dx.doi.org/10.1016/j.rmed.2010.05.014>
-
58. Washko GR, Lynch DA, Matsuoka S, et al. Identification of early interstitial lung disease in smokers from the COPD Gene Study. *Acad Radiol.* 2010;17(1):48-53.
doi: <http://dx.doi.org/10.1016/j.acra.2009.07.016>
-
59. Washko GR, Hunninghake GM, Fernandez IE, et al. Lung volumes and emphysema in smokers with interstitial lung abnormalities. *New Eng J Med.* 2011;364:897-906.
doi: <http://dx.doi.org/10.1056/NEJMoa1007285>
-
60. Doyle TJ, Washko GR, Fernandez IE, et al. Interstitial lung abnormalities and reduced exercise capacity. *Am J Crit Care Med.* 185(7):756-762.
doi: <http://dx.doi.org/10.1164/rccm.201109-1618OC>
-
61. Hunninghake GM, Hatabu H, Okajima Y, et al. MUC5B promoter polymorphism and interstitial lung abnormalities. *New Eng J Med.* 2013;368:2192-2200.
doi: <http://dx.doi.org/10.1056/NEJMoa1216076>